

The anisotropy of inverse beta decay and antineutrino detection

G.Fiorentini, M.Moretti, F.L.Villante

Dipartimento di Fisica, Universita' di Ferrara, I-44100, Ferrara (Italy)
Istituto Nazionale di Fisica Nucleare, Sez. di Ferrara, I-44100, Ferrara (Italy)

The anisotropy of the positrons emitted in the reaction $\bar{\nu}_e + p \rightarrow n + e^+$ has to be taken into account for extracting an antineutrino signal in Superkamiokande. For the Sun, this effect allows a sensitivity to $\nu_e \rightarrow \bar{\nu}_e$ transition probability at the 3% level already with the statistics collected in the first hundred days. For a supernova in the Galaxy, the effect is crucial for extracting the correct ratio of $\nu - e$ to $\bar{\nu}_e - p$ events.

As well known, the specific signature of antineutrinos in hydrogen containing materials is through the inverse beta decay (I β D), $\bar{\nu}_e + p \rightarrow n + e^+$, which produces *almost isotropically* distributed monoenergetic positrons ($E_{e^+} = E_{\bar{\nu}} - \Delta m$; $\Delta m = m_n - m_p$). For energy above a few MeV, the differential cross section is:

$$\frac{d\sigma}{d\cos\theta} = \frac{\sigma_0(E_{\bar{\nu}})}{2}(1 - a\cos\theta) \quad (1)$$

where:

$$a = \frac{(g_A/g_V)^2 - 1}{3(g_A/g_V)^2 + 1} \simeq 0.1 \quad , \quad (2)$$

g_V (g_A) is the vector (axial) coupling of the neutron and $\sigma_0(E_{\bar{\nu}})$ is the total cross section for antineutrino energy $E_{\bar{\nu}}$.

The goal of this paper is to assess the importance of the angular dependence of the cross section in two situations of practical interest:

- The search for solar antineutrinos;¹
- the detection of a supernova in the Galaxy.

1. Solar Antineutrinos

In the absence of a solar antineutrino flux, the Superkamiokande (SK) background is expected to be isotropic. In the presence of solar antineutrinos, positrons emitted by I β D contribute to the

¹This signal is of interest since it might be a signature of ν decay or magnetic moment. see [1] for details and references

background, which, due to the angular dependence of I β D cross section, should have a non-zero angular slope proportional to the antineutrino flux. A linear fit to the counting yield, $C = C_0 - C_1 \cos\theta$ (in the angular region where events from the $\nu - e$ interactions can be neglected, see Fig. 1), provides the antineutrino flux $\Phi_{\bar{\nu}}$ through the relation:

$$\Phi_{\bar{\nu}}(E_{\bar{\nu}} > E_0) = \frac{C_1}{T} \frac{2}{N_p \epsilon \bar{\sigma}_0 a} \quad (3)$$

where N_p is the number of free protons, T is the exposure time, ϵ is the (assumed constant) detection efficiency, E_0 the minimal detectable antineutrino energy and $\bar{\sigma}_0$ is the cross section averaged over the antineutrino spectrum for $E_{\bar{\nu}} > E_0$.

In order to provide a quantitative illustration of the previous points we used data from the first 101.9 operational days of SK, as reported in fig.3 of [2], corresponding to $E_0 = 8.3\text{MeV}$. The reported background does not show an angular dependence. According to equation (3), to extract an upper limit on the solar antineutrino flux, we must know the average cross section $\bar{\sigma}_0$, which can be determined within two approaches:

a) Assuming that the antineutrino spectrum has the same shape as that of ^8B solar neutrinos, one has $\bar{\sigma}_0 = 7.1 \cdot 10^{-42} \text{cm}^2$. This gives, as a final result, $\Phi_{\bar{\nu}}(E_{\bar{\nu}} > 8.3\text{MeV}) < 6 \cdot 10^4 \text{cm}^{-2} \text{s}^{-1}$, to the 95% C.L. This bound corresponds to a fraction $x=3.5\%$ of the solar neutrino flux (in the energy range $E_{\nu} > 8.3\text{MeV}$) predicted by the SSM [3].

b) As σ_0 is an increasing function of $E_{\bar{\nu}}$, one has $\bar{\sigma}_0 \geq \sigma_0(E_0) = 4.5 \cdot 10^{-42} \text{cm}^2$. This lower limit




Figure 1. Sketch of the expected angular distribution of events in the presence of a solar $\bar{\nu}_e$ flux.

to $\bar{\sigma}_0$ gives a *model independent* bound $\Phi_{\bar{\nu}}(E_{\bar{\nu}} > 8.3\text{MeV}) < 9 \cdot 10^4 \text{cm}^{-2}\text{s}^{-1}$ to the 95% C.L.

We remark two points of the method just presented:

- The sensitivity to antineutrinos increases as statistics accumulates. Within three years of data taking, the sensitivity to $\nu_e \rightarrow \bar{\nu}_e$ transition probability will reach the 1% level, thus allowing for a definite test of several theoretical models.
- The determination of the angular slope C_1 provides a mean for *detecting* antineutrinos from the Sun (and not only for deriving upper bounds). A non vanishing slope for SK background would be, in fact, a clear signature of a solar antineutrino flux.

2. Supernova detection

If a type-II supernova explosion occurs near the center of our galaxy about 4000 $\bar{\nu}_e + p \rightarrow n + e^+$ events and 300 $\nu - e^- \rightarrow \nu - e^-$ events will be produced in the Superkamiokande detector [4]. The

detection of supernova neutrinos and antineutrinos will provide important information on the supernova mechanism. In particular we remark that:

- from the strong correlation of the elastic scattering events one can reconstruct the supernova direction with an accuracy of about four degrees [4];
- from a fit to experimental data one can determine the number of neutrinos and antineutrinos events, thus providing important inputs to check supernova modelling.

The anisotropy of $I\beta D$ is so small that it looks reasonable to neglect it when discussing the detector sensitivity [4]. However, let us investigate how it can affect the data analysis. With this aim we have simulated a supernova explosion in the galactic plane, assuming that it produces $N_{e+} = 4000$ and $N_{e-} = 300$ events in the detector. We took into account a 30° angular resolution [4] and repeated the simulation two-hundred times. We analysed the data looking for the galactic longitude ϕ of the supernova and the ratio $f = N_{e-}/N_{e+}$ of reconstructed events, see fig.2.

If the analysis is performed neglecting the anisotropy (i.e. $a = 0$) the supernova direction is correctly determined within the predicted accuracy, however the reconstructed f is badly underestimated ($f = 0.057 \pm 0.009$, the correct value being $f = 0.075$) and the fit is poor ($\chi^2/DOF = 24/18$ on the average of the 200 simulations, using 20 angular bin). We checked that when the anisotropy is included, both the supernova direction and the ratio f are correctly reconstructed, and the quality of the fit improves significantly. These features are clearly understood:

- as the positron angular distribution is symmetrical around the supernova direction, it does not alter its reconstruction.
- The depleted forward positron production mimics a reduced number of $\nu - e$ events.

In conclusion, we remark that inclusion of the anisotropy of $I\beta D$ is essential for extracting the correct ratio of $\nu - e$ to $\bar{\nu}_e - p$ events.




Figure 2. The simulated counts as a function of the galactic longitude for a supernova at the galactic center (dots), with their statistical errors (bars), the best fit for $a = 0$ (dotted curve) and that for $a = 0.1$ (full curve)

REFERENCES

1. G. Fiorentini, M. Moretti, F.L. Villante, Phys. Lett. B, in press, 1997
2. Y. Totsuka, "First result from Super-Kamiokande", presented at Texas Symposium (1996).
3. J.N. Bahcall and M.H. Pinsonneault, Rev. Mod. Phys. 61 (1992) 885.
4. K. Nakamura, "Superkamiokande", presented at 3rd NESTOR International Workshop, Pylos, Greece, 1993